FAQ: Sorption and Phosphorus Stability in Utah Lake

What is sorption, and why does it matter in Utah Lake?

Sorption is the process by which dissolved substances—like phosphorus (P)—attach (and detach) to solid surfaces, such as lakebed sediments. In Utah Lake, sorption plays a dominant role in regulating phosphorus levels. The lake's shallow depth, high sediment mixing, and phosphorus-rich sediments allow for rapid and reversible exchange between the water column and sediments, buffering P concentrations and keeping them stable.



Why is Utah Lake especially prone to sorption-dominated phosphorus cycling?

Utah Lake is well suited for sorption processes to play a large role:

- It is shallow (average depth ~3 meters), well-mixed due to wind and wave action, and doesn't have a thermocline.
- It has phosphorus-rich sediments from natural geologic sources like the Phosphoria and Park City formations.
- It experiences frequent sediment resuspension due to wind and carp bioturbation.
- Sediment phosphorus concentrations range from 300–1100 mg/kg, with ~40% being available for exchange with the water column.

Does the phosphorus concentration in Utah Lake fluctuate with lake volume?

Surprisingly, no. Long-term monitoring (1989–2023) shows that dissolved phosphorus (P) concentrations remain stable, typically between 0.02 and 0.04 mg/L, even though lake volume can annually vary by more than 50% due to snowmelt, runoff, and evaporation with year-to-year variations by a factor of 2. This defies standard mass-balance expectations and reinforces that sorption dominates the system rather than P inflows or outflows (i.e., mass balance).

What is a K_d Value and Why Does It Matter?

The **K_d value**, or *distribution coefficient*, is a measure of how strongly phosphorus (P) binds to sediments versus staying dissolved in water. A high K_d means phosphorus tends to **stick to sediments** (strong sorption with the concentration in the sediment higher than that in the water), while a low K_d means more phosphorus stays in the water (the concentration in the water is higher than the sediment). In Utah Lake, K_d values range from **hundreds to over 1000**, depending on conditions like pH and sediment composition. These high values indicate that **most dissolved phosphorus quickly binds to sediments**, which helps explain why phosphorus concentrations in the water remain stable—even when external phosphorus inputs or lake volumes change.



What would we expect without sorption?

In a typical lake without strong sorption or where the sediments do not interact as much with the water column:

- P concentrations would increase in dry years or during late summer when the lake shrinks through evaporation, but external loads remain similar.
- P concentrations would decrease in wet years due to dilution from spring runoff or precipitation (again while external loads remain similar).

However, Utah Lake's data show the opposite: P concentrations stay nearly constant, suggesting buffering from sorption on the sediment.

How does sorption respond to phosphorus inflow or loading?

Sorption buffers against external phosphorus loads—like wastewater or stormwater—increasing water concentrations. Even large changes in phosphorus input have little effect on the lake's water column:

- Raising P from 0.030 to 0.035 mg/L would require adding 346–714 metric tons of phosphorus.
- To reduce P from 0.030 to 0.025 mg/L, Utah Lake would need to be flushed 13–27 times over with clean water (depending on the K_d value—13 times is for a K_d value of 100, the lower end of the range).

This level of intervention far exceeds current or practical management options.

Why doesn't reducing P loads to Utah Lake lower water column concentrations?

Because the sediments contain an enormous reservoir of phosphorus, and sorption seeks equilibrium between sediment and water. Reducing external P input does not change that equilibrium quickly because of the large amount of P in the sediment that can "make up the difference". Instead, sediment P continues to buffer water column P concentrations, resisting change. This makes Utah Lake insensitive to typical load-reduction strategies.



What are the policy implications of a sorption-dominated lake?

Since reducing external phosphorus loads may not significantly lower water column concentrations, management efforts may be more effective if they focus on:

- Sediment stabilization (e.g., carp removal, vegetation restoration) to reduce sediment-water mixing.
- In-lake treatments targeting internal P cycling.
- Improving predictive models that account for sorption dynamics to provide Managers with realistic options for consideration.

Is this situation unique to Utah Lake?

Not entirely. Other shallow, well-mixed lakes with P-rich sediments—especially those underlain by geologic phosphate—may exhibit similar behavior. However, Utah Lake's combination of shallow depth, extensive sediment mixing, and high sediment P concentrations from geologic P sources makes it a sorption-dominated lake.

For more information:

Taggart et al. (2024). *Historical Phosphorus Mass and Concentrations in Utah Lake*. Water 16(933).

Jarvis (2023). *Phosphorus Sorption and Kinetics Across the Sediment-Water Interface of Utah Lake*. Thesis, BYU ScholarsArchive.

Aremu (2023). *Investigating the Geochemical Trends at the Sediment-Water Interface of the Shallow and Eutrophic Utah Lake.* Thesis, BYU ScholarsArchive.


water
MDPI

Article

Historical Phosphorus Mass and Concentrations in Utah Lake: A Case Study with Implications for Nutrient Load Management in a Sorption-Dominated Shallow Lake

Jacob B. Taggart [1], Rebecca L. Ryan [1], Gustavious P. Williams [1,*], A. Woodruff Miller [1], Rachel A. Valek [1], Kaylee B. Tanner [1] and Anna C. Cardall [2]